

Grain Sorghum **Weed Control Guide**



Contents

Current farming practice	4
Current herbicide strategies in grain sorghum	5
Key weeds by regions	6
Visual weed identification – key weeds	7
Herbicides	10
Weed control guide	12
Weed suppression by growing a competitive sorghum crop	19
Introduction to igrowth sorghum	24
Good practices for the management of tolerant crops to herbicides	28
Integrated weed management and stewardship	29
How do herbicide resistance weeds develop?	29

List of tables

Table 1: Key weeds by region – Central Queensland	6
Table 2: Key weeds by region – North East & QLD South East	6
Table 3: Key weeds by region – NSW North West & QLD South West	6
Table 4: Registered herbicides	10
Table 5: Grass weeds – Pre-emergence	12
Table 6: Broadleaf weeds – Pre-emergence	12
Table 7: Weeds – Post-emergence	13
Table 8: Re-crop intervals required for herbicides that may be utilised prior to sorghum planting	17
Table 9: Susceptibility of grass weeds to IMI herbicides	25
Table 10: Susceptibility of broadleaf weeds to IMI herbicides	25
Table 11: Re-crop intervals from in-crop applications of Intervix, Lightning and Spinnaker or fallow applications of Flame in summer rainfall areas?	27



Current farming practice

Dryland farming

Dryland grain sorghum production accounts for the largest proportion of Sorghum area under cultivation where crop production is underpinned by the accumulation of stored soil water in the fallow preceding planting. In some instances, sorghum has been grown as a “double-crop” after winter crop harvest where sufficiently rainfall is received around or soon after the harvest period to provide at least some replenishment of subsoil moisture and sufficient planting moisture to justify a planting decision.

In northern New South Wales (NSW) outside of the higher rainfall areas, dryland sorghum is more often than not grown on a longer fallow from a previous winter crop with spring or early summer planting preferred depending on the timing of planting rains.

In more favoured areas and further north where summer growing season rainfall is usually greater, rotations tend to be shorter with sorghum grown following either sorghum or another summer crop including dryland cotton, occasionally sunflower or pulse crop such as mungbean.

Chickpea is frequently planted in rotation with sorghum in southern Queensland (QLD) where a double crop makes up a significant proportion of area of chickpea planted on the central and eastern Darling Downs regions.

Irrigated farming

Areas of irrigated production vary with availability of irrigation water and the relative profitability of alternative crops. In recent decades, irrigated cotton has comprised the largest footprint of broadacre crops displacing traditional grain crops in some areas. Sorghum has been no exception to this though there are still significant areas of sorghum grown under irrigation and in particular northern NSW and southern QLD. Irrigated crops make up the greater proportion of crops grown in central and southern NSW where the incidence of growing season rainfall makes dryland production unreliable.

Traditional production systems have utilised a flood/furrow irrigation layout though increasingly, overhead irrigation systems delivering water via centre pivot or lateral move systems have accounted for significant areas of production.

Geographical distribution of grain sorghum

Grain sorghum is grown over a wide latitude extending as far south as the Riverina district in southern NSW to the northern central highlands in Queensland. Sorghum has also been grown outside this area over the history of production but currently, no extensive areas for grain production are consistently planted outside this general area.




Figure 1: Northern NSW and Southern QLD grain sorghum growing areas

Current herbicide strategies

In-crop weed control is largely borne by the use of atrazine and metolachlor/s-metolachlor as pre-emergence treatments and atrazine and fluroxypyr as post-emergence herbicides. Interrow cultivation is still extensively used in flood/furrow systems but the use of interrow cultivation has declined over time for dryland production where zero and minimum tillage farming systems have displaced conventional farming practices.

Other selective pre-emergence herbicides include terbuthylazine (Terbyne) while this product is also used as a substitute for the lower-priced atrazine as a post-emergence herbicide in partner with fluroxypyr. Products containing Picloram and 24D are used less due to the superior crop safety profile of fluroxypyr plus atrazine/terbuthylazine. Bromoxynil, while registered for use, is rarely if ever used. Triclopyr e.g. Garlon 600 is occasionally used in post-emergence mixtures to enhance control of prickly paddy or camel melon.

Valor (flumioxazin) has been utilised in fallow leading up to various crops as a residual herbicide and provides an important tool in providing pre-emergence control of Feathertop Rhodes grass and sowthistle in particular.

Interrow shielded application of glyphosate has been used in the past but this never became a widely practiced method of weed control except in salvage situations where pre-emergence herbicides were not applied, and grass-weed control could not be affected using selective post-emergence herbicides.



Shielded Sprayer. Photo courtesy of Hayes Spraying

Post-emergence options for grass-weed control remains an outstanding vulnerability of weed control in grain sorghum, this almost completely being reliant on s-metolachlor/metolachlor and to a much lesser extent atrazine and terbuthylazine.

Triazine herbicides and in particular atrazine has declined in effectiveness against grass weeds through a combination of evolving resistance of species such as awnless barnyard grass but also commonly through a shift toward incidence of grass species having a low or inconsistent susceptibility to triazine herbicides. These species include liverseed (urochloa) grass, Feathertop Rhodes grass, sweet summer grass (central Qld), sorghum spp., dinebra grass, crowsfoot grass, stink (black) grass, native millet, spring grasses, windmill grass, button grass and a number of other native perennial grasses which have become more prominent in zero-tillage production systems.

Key weeds by region

Table 1: Central Queensland

RANK	WEED	AREA (HA)
1	Feathertop Rhodes grass	31,313
2	Sweet summer grass	24,093
3	Parthenium weed	2,171
4	Bellvine	1,661
5	Fleabane	1,571
6	Wild mustard	345
7	Barnyard grass	329

Table 2: New South Wales - North East, and Queensland - South East

RANK	WEED	AREA (HA)
1	Barnyard grass	80,264
2	Fleabane	62,868
3	Feathertop Rhodes grass	37,649
4	Sweet summer grass	31,827
5	Bathurst burr	23,969
6	Noogoora burr	10,538
7	Wild oats	4,108
8	Thornapple	3,286
9	Yellowvine/Caltrop	3,072
10	Wireweed	2,143

Table 3: New South Wales - North West, and Queensland - South West

RANK	WEED	AREA (HA)
1	Barnyard grass	6,496
2	Feathertop Rhodes grass	4,452
3	Fleabane	405
4	Melons	405
5	Sowthistle	405
6	Noogoora burr	405

Source: Llewellyn RS, Ronning D, Ouzman J, Walker S, Mayfield A and Clarke M (2016) *Impact of Weeds on Australian Grain Production: the cost of weeds to Australian grain growers and the adoption of weed management and tillage practices*

Report for GRDC. CSIRO, Australia.

Visual weed identification – key weeds



Awnless barnyard grass *Echinochloa colona*



Liverseed (urochloa) grass *Urochloa panicoides*



Sowthistle *Sonchus oleraceus*



Feathertop Rhodes grass *Chloris virgata*



Red root amaranth (Redshank) *Amaranthus hybridus*



Bathurst burr *Xanthium spinosum*

Visual weed identification – key weeds



Crowsfoot grass *Eleusine indica*



Rhynchosia *Rhynchosia minima*



Red pigweed *Portulaca oleracea*



Black pigweed *Trianthema portulacastrum*



Caltrop *Tribulus terrestris*



Flax-leaf fleabane *Conyza bonariensis*



Cowvine (peachvine) *Ipomea lonchophylla*



Blackberry nightshade *Solanum nigrum*



Bladder ketmia *Hibiscus trionum* ssp. *Trionum*



Dwarf amaranth *Amaranthus macrocarpus*



Yellowvine *Tribulus micrococcus*

Herbicides

Table 4: Registered herbicides

HERBICIDE ACTIVE	CONCENTRATION	EXAMPLE TRADENAMES	APPLICATION TIMING	RATES
Atrazine	600 g/L SC	Gesaprim 600	Pre & Post 4 - 6 leaf	
	900 g/kg WG	Gesaprim 900	Pre & Post 4 - 6 leaf	2 kg (PRE); 2.5 kg (POST) or 830-1100 g/ha + fluroxypyr (POST)
Terbuthylazine	850 g/kg WG	Terbyne Xtreme	Pre & Post 5 - 7 leaf	1.2 kg/ha or 0.95 kg/ha + fluroxypyr
s-metolachlor	960 g/L EC	Dual Gold, Bouncer 960S	Pre	1 - 2 L/ha
Metolachlor	720 g/L or 960 g/L EC	Various	Pre	2 - 4 L/ha (720EC) 1.5 - 3 L/ha (960)
Fluroxypyr	200 g/L EC	Flagship 200	Post 4 leaf - boot	500 - 1500 mL/ha
	333 g/L EC	Starane Advanced	Post 4 leaf - boot	300 - 900 mL/ha
	400 g/L EC	Comet 400, Flagship 400	Post 4 leaf - boot	250 - 750 mL/ha
Triclopyr	600 g/L EC	Garlon 600	Post 4 - 6 leaf	80 mL/ha
Bromoxynil	200 g/L EC	Bromicide	Post 4 - 12 leaf	1500 - 2000 mL/ha
	400 g/L EC	Bronco 400	Post 4 - 12 leaf	750 - 1000 mL/ha
Picloram + 2,4-D		Enforcer 75D	Post 4 - 6 leaf	330 or 500 mL/ha
Propachlor	480 g/L SC	Ramrod	Pre	11 - 13.5 L/ha
Halosulfuron	750 g/kg	Sempra	Post Up to 60 cm	65 to 130 g/ha
2,4-D	Various	Not all brands registered	Post 4 - 8 leaf	
Dicamba	500 g/L SL	Not all brands registered	Post 3 - 6 leaf	280 - 320 mL/ha
Imazamox + imazapyr	48 g/L SL	Intervix	Post 2 - 6 leaf (IG sorghum only)	750 - 1000 mL/ha + Hasten or Kwickin
Glyphosate	Various	Roundup Crucial Wipeout	Post shielded	
Diquat	Various	Reglone	Pre-harvest	2000 - 3000 mL/ha

Comments on herbicides

- Dicamba and 2,4-D are rarely used as these can cause significant crop injury when applied over-the-top.
- Not all brands of 2,4-D are registered as risks of crop injury are considered to outweigh commercial risk to the registration holder.
- Picloram + 2,4-D is rarely used due to marginal crop selectivity in some situations.
- Halosulfuron can cause crop shortening and stunting particularly at the higher rate.
- Propachlor is rarely if ever used due to high cost when compared to alternative treatments.
- Atrazine is compatible with s-metolachlor, metalochlor, propachlor, fluroxypyr, triclopyr, bromoxynil, picloram + 2,4-D, dicamba and 2,4-D.
- Metolachlor and s-metolachlor are registered for use prior to, at or immediately following planting. A label extension for use post-emergence has been granted.
- Non-ionic surfactant is recommended for post-emergence applications including atrazine.
- An esterified vegetable oil surfactant such as Hasten is recommended for post-emergence applications of terbutylazine or Intervix.



Examples of 24D damage in Grain Sorghum

Weed control guide

Table 5: Grass weeds – Pre-emergence

WEED	ATRAZINE	METOLACHLOR S-METOLACHLOR	TERBUTHYLAZINE E.G. TERBYNE XTREME
Barnyard grass	x§	x	x§
Liverseed (urochloa) grass	o	x	o
Feathertop Rhodes grass	o	x	o
Sweet summer grass	o	x	o
Stink (black) grass	x	x	x
Summer grass	x	x	x
Crowsfoot grass	x	x	x
Spring grass	x	x	x
Native millet	o	x	o

x Controlled
 o Not controlled
 § Control is unsatisfactory where there has been a long history of use due to evolution of resistant populations.

Table 6: Broadleaf weeds – Pre-emergence

WEED	ATRAZINE	METOLACHLOR S-METOLACHLOR	TERBUTHYLAZINE
Amaranth	x	x	x
Annual ground cherry	x	o	x
Bellvine/cowvine	o	o	o
Blackberry nightshade	x	x¶	x
Bladder ketmia	x	o	x
Burrs	x¶	o	x¶
Caltrop/yellowvine	x¶	o	x¶
Common thornapple	x¶	o	x¶
Dwarf amaranth	x	x	x
Fleabane	x	o	x
Melons	x¶	o	x¶
Mintweed	x	x	x
Parthenium weed	x	o	x
Pigweed (black)	x	x	x
Pigweed (red)	x	x	x
Sowthistle (milk thistle)	x¶	o	x
Wireweed	x	o	x

x Controlled
 o Not controlled
 ¶ Variable control

Table 7: Post-emergence

RANK	ATRAZINE	TERBUTHYLAZINE E.G. TERBYNE XTREME	FLUROXYPYR E.G. STARANE, COMET, FLAGSHIP	TRICLOPYR E.G. GARLON 600	FLUROXYPYR + ATRAZINE	FLUROXYPYR + TERBU-THYLAZINE	PICLORAM + 2,4-D PLUS ATRAZINE	BROMOXNYL E.G. BROMICIDE + ATRAZINE	INTERVIX
Broadleaf weeds									
Amaranth	x	x	o		x	x	x	o	x
Annual ground cherry			x		x	x	x	x	x
Bellvine/cowvine					x	x	x	x	x
Blackberry nightshade			x	x	x	x			s
Bladder ketmia	x	x	o		x	x	x	x	s
Burrs		x	x		x	x	x		x
Caltrop/ yellow-vine		x	o		x	x		x	x
Common thornapple	x	x	x		x	x	x	x	x
Dwarf amaranth	x	x	o		x	x	x	x	x
Fleabane			o		x		x		o
Melons			o	x	x	x	x		s
Mintweed	x	x	o		x	x	x	x	s
Parthenium weed	x		o		x	x	x		
Pigweed (black)	x	x	o		x	x	x	x	s
Pigweed (red)			x		x	x	x		x
Sowthistle (milk thistle)	o	x	x		x	x	x		s
Wild gooseberry		x							x
Wireweed			o						s
Grass weeds									
Barnyard grasses	s	o	o	o	o	o	o	o	x
Crowsfoot grass	o	o	o	o	o	o	o	o	s
Liverseed grass	o	o	o	o	o	o	o	o	x
Feathertop Rhodes grass	o	o	o	o	o	o	o	o	o
Johnson grass	o	o	o	o	o	o	o	o	s
Summer grass	o	o	o	o	o	o	o	o	s

x Controlled
o Not controlled

s Variable control
s Suppression

Use of Residuals


Reliance on pre-emergence herbicides for grass weed control has been a vital component for weed control in sorghum due to the lack of effective and selective post-emergence herbicide options, other than with shielded applications with glyphosate.

Initially, atrazine provided effective pre and post-emergence control of a wide range of weeds. However, a number of grass species were either not effectively controlled including liverseed grass (some populations were partially susceptible initially) and many native grasses including native millet, Qld blue grass and spring grasses proved tolerant. Further, as use of atrazine became regular in sorghum production, effectiveness appeared to decline on important grass species such as awnless barnyard grass due possibly to evolution of resistant weeds and/or the accelerated bio-degradation of atrazine which occurs when soil microflora capable of degrading atrazine in the soil, build up in number as use intensifies.

Useful Link:

ICAN Herbicide and Weed manuals: www.icanrural.com.au/resources.html

Figure 2: Sorghum growth stages by herbicide applications



Pre-sow/
pre-emergent
knock down
and residual
top up

In-crop knock down
and residual

In-crop knock
down and residual

Harvest
aid /
late
knock-
down

Early fallow
application of
pre-emergent
Grass control
products

Source: Adapted from NSW DPI Grain Sorghum Ag Facts.

Broadleaf options

Many broadleaf weeds are consistently and well controlled by pre-emergence applications of either atrazine or terbuthylazine where these germinate on rainfall or over-head irrigation following application. Inconsistency in control is often the result of weeds having germinated prior to application but not controlled at the point of treatment. This is frequently the situation where treatments are applied at planting where the planting rain initiates germination, but these fail to emerge in time for knockdown sprays to be able to remove them prior to crop emergence.

A further problem may occur where hard-to-kill weeds such as flax-leaf fleabane are not effectively controlled at the point of planting due either to difficulties in achieving good coverage in standing stubble, or due to restriction in options available for knockdown of well-established weeds at the point of planting.

A good option to avoid either of these situations, is to consider bringing forth application of pre-emergence herbicides prior to a planting rain or in the case of fleabane even earlier where the onset of spring germination can occur from early August once soil temperatures rise and spring rainfall stimulate emergence.

Pre-emergence control of deep germinating broadleaf weeds is also problematic with pre-emergence herbicides where dryness in the surface layers and moisture below can facilitate germination of weeds in the region of soil where herbicide is not present. Weed species where this is observed include cowvine (peachvine), bellvine, burrs (Bathurst and Noogoora), thornapples, bladder ketmia and native legumes including Rhynchosia, native sensitive weed, native vigna. In addition, pre-emergence control of yellowvine or caltrop also proves problematic with atrazine and terbuthylazine though under wet conditions, control is more effective.

Post-emergence application of herbicides is frequently a better strategy where the weed spectrum is made up chiefly of weeds that germinate from depth or include native legumes. In this case, combinations of either atrazine or terbuthylazine with fluroxypyr (e.g. Starane Advanced, Comet 400, Flagship 400) represents a more cost-effective approach. Where weed numbers are expected to be high and where other more susceptible weeds to pre-emergence herbicides are present, a pre-emergence application either in the fallow prior to sorghum or at the point of planting remains as a sound option.

Grass options

Pre-emergence treatments for control of grasses depends very much on the timeliness of application of s-metolachlor or metolachlor in relation to the expected emergence time for grasses. Like some broadleaf weeds that begin to emerge during late winter and early spring, grass control can be compromised if applications of pre-emergence treatments take place following the initiation of germination of grass weeds particularly where this takes place simultaneous with crop planting and herbicide application.

In the case of Feathertop Rhodes grass, germination can take place virtually at any time except under very cold conditions which means that when temperatures are sufficiently high for sorghum planting, Feathertop Rhodes may have already germinated and options for knockdown control are virtually exhausted where populations exhibit tolerance to glyphosate sprays.

A strategy recently employed on some farms is to bring forward the application of s-metolachlor or metolachlor well in advance of the first significant emergence of Feathertop Rhodes grass. In southern Qld, this requires applications any time from early to mid-August prior to rainfall events of at least 10 mm on medium clays or 15 to 20 mm on heavy clay soils.

While it is preferred to time applications within a couple of weeks of application, treatments can remain effective for some time without the occurrence of activating rainfall though expected duration of effect of metolachlor will decline in the absence of rainfall over time.

Mechanical incorporation after application is a viable strategy where the field has no stubble or is conventionally formed. Mechanical incorporation should insure the chemical is incorporated to a depth greater than 1-2cm.

For other spring germinating grasses such as barnyard grass, applications can be delayed until late September or early October as the emergence period for this weed does not commence as early as for Feathertop Rhodes. Pre-emergence control of liverseed (urochloa) grass can be problematic as in many years, emergence can take place during winter months if there are sufficiently warm temperatures around the time of rainfall events from late autumn. In this instance, it is better to deal with any early emergence with cultivation or knockdown spray and apply pre-emergence treatments to a "brown field" or at least at a time when there is no germination taking place. In instances where fallow applications of metolachlor take place prior to planting rainfall, it is recommended that a "top up" be applied following planting.

As a guide, early application should be around the equivalent of 1.5 L/ha Dual Gold (1440 ga/ha s-metolachlor) though this could be reduced to around 1.0 L/ha if a planting rain is imminent. Follow-up applications post-planting pre-emergence should start at 0.5 L/ha and increase to 1.5 L/ha if an extended period has occurred following the initial fallow application.

Useful Link:

Keeping Sorghum Safe when using metolachlor based herbicides: <https://www.grdc.com.au/TT-KeepingSorghumSafe>

Improving effectiveness and safety of Dual Gold and other metolachlor herbicides

Split applications of s-metolachlor and metolachlor not only benefit with the effectiveness of weed control but also mitigate risks associated with the application of a single application of a higher rate at the point of planting.

Crop injury has been associated with moderate to heavy rainfall events soon after application where higher rates of metolachlor have been applied. By splitting application between a fallow application in advance of planting rain and topping up with low to moderate rates (0.5 - 1.0 L/ha Dual Gold) at planting, risks of crop injury are reduced.

The critical time for injury from metolachlor appears to be within 0 and 5-7 days of planting at a time when germination is occurring. Damage is also more common for plants germinating closer to the soil surface or where the effect of heavy rainfall or overland water flow is to reduce the effective depth of crop germination to bring germinating seeds within 2 - 4 cm of the soil surface. Planting with low vigour seed and occurrence of low soil temperatures are also risk factors for metolachlor injury as this increases the period of vulnerability of exposure to metolachlor and reduces the ability of plants to withstand injury relative to seed batches with strong vigour or where soil temperatures facilitate more rapid emergence.

Alternative pre-emergence grass control options in fallow

Valor® (flumioxazin) has been recently utilised for pre-emergence weed control in fallow at rates of 210 - 280 g/ha providing an alternate mode of action to s-metolachlor for the control of Feathertop Rhodes grass in particular. Unlike s-metolachlor, Valor needs to be applied at least one month prior to planting – a use pattern that lends itself to winter application prior to first germination of Feathertop Rhodes grass.

The low solubility of Valor means that weed seeds germinating close to the surface are well controlled including broadleaf weeds such as sowthistle and fleabane. Deeper germinating weeds such as liverseed grass are less reliably controlled.

Re-crop intervals following use of various herbicides in prior crops or fallow

Intervals to re-cropping to sorghum can be an important consideration particularly when utilising herbicides in fallow and sometimes in prior crops where a shorter fallow interval occurs. In addition, herbicides that nominally have no useful pre-emergence activity but are used in fallow closer to planting time require careful use to minimise possibility of damage to sorghum.

Table 8: Re-crop intervals required for herbicides that may be utilised prior to sorghum planting

HERBICIDE	TRADENAME	INTERVAL	RAINFALL REQUIREMENT	COMMENTS
2,4-D	Several	3, 7 or 10 days	Nil	Interval increases with increasing rate
2,4-D + picloram	Tordon Fallowboss	4 months	100 mm	Rates up to 700 mL/ha
Atrazine	Several	0		
Clopyralid	Lontrel	7 or 14 days	50 mm	
Dicamba	Several	1, 3 or 7 days		Interval increases with increasing rate. 1 day in Qld
Flumioxazin	Valor	1 month (210-280g)		Zero interval for 30 or 45 g/ha
Fluroxypyr	Starane	7 days		
Halauxifen + aminopyralid	Trezac	3 months	100 mm	
Halauxifen + florasulam	Paradigm	4 months	100 mm	
Halauxifen + fluroxypyr	Pixxaro	1 month	100 mm	
Imazapic	Flame	10 months		Shorter for IG sorghum
Haloxfop	Verdict	12 weeks		
Isoxaflutole	Balance	7 months	250 mm	
s-metolachlor	Dual Gold	0		
Metsulfuron	Ally	14 months		
Propaquizafop	Shogun	28 days		
Saflufenacil	Sharpen	1 day		
Terbuthylazine	Terbyne	0		
Tribenuron	Express	7 days		21 days where soil temperatures <15 C

Pre-harvest glyphosate as harvest aid

Pre-harvest application of glyphosate for control of regrowth and facilitation of a more efficient herbicide was a technique first developed in the mid 1980's and has become standard practice over much of the sorghum growing regions of Australia.

As an indeterminant plant, sorghum harvest is frequently frustrated by the ongoing production of growth particularly where late season rainfall or irrigation allows for continuing growth. Some varieties have a strongly perennial habit with the result that harvest can be delayed for long periods while immature tillers develop to bear seed heads that eventually will produce viable grain or be terminated by killing frosts. Timely application of glyphosate sprays will prevent or suppress growth of late-season tillers allowing for more efficient and timely harvest. Desiccation of leaf matter and in some instances, acceleration of grain dry-down allow for more efficient and clean harvesting operation.

Early termination of growth also increases opportunity for moisture conservation in the soil profile, maximising the opportunity for double cropping in the event there is sufficient replenishment of soil moisture reserves before the window of winter cropping closes.

Timing of application is critical for the success of pre-harvest applications of glyphosate. Most consistent and effective results are achieved where late immature tillers are not present at the time of application and sprays are applied at physiological maturity of primary tillers. This can be judged without the aid of moisture meters with the appearance of a 'black layer' at the base of individual seeds in the seed head.

As a guide, grain moisture below 30 per cent is considered optimal for timing of pre-harvest sprays. Applications made prior to physiological maturity of primary tillers are less likely to be successful and may result in delayed maturity as plants attempt to complete grain fill. Early application can also lead to an extended interval to harvestable grain moisture leaving plants open to lodging should conditions deteriorate due to untimely rainfall.

In the event of late season rainfall, it is better to apply treatments before the initiation of late tillers or at least before there is opportunity for late tillers to develop immature seed-heads.

The advent of high clearance ground-sprayers has enabled more efficient and timely control of sorghum with pre-harvest applications. While aerial application has been successfully used in the past, uniformity of spray deposit may be compromised by the need for a higher release height above the crop canopy, and the interference with deposition by a relatively dense crop canopy and varying wind velocity at height.

Best results with ground application are achieved with applications made in at least 50 L/ha carrier volume.

A medium to coarse spray quality is likely to be equally effective in achieving satisfactory results while for aerial application, a minimum spray volume of around 30 L/ha is optimal for achieving more uniform deposition using narrower swath widths than would normally consider necessary for fallow spraying.

Useful Link:

GRDC Grownotes Sorghum Section 11 Crop Desiccation / Spray Out:

https://grdc.com.au/__data/assets/pdf_file/0018/370602/GrowNote-Sorghum-North-11-Desiccation.pdf

Weed suppression by growing a competitive sorghum crop

Michael Widderick QDAF

Summer grass weeds awnless barnyard grass (BYG) (*Echinochloa colona*) and feathertop Rhodes grass (FTR) (*Chloris virgata*) are problematic in sorghum crops. Not only are effective herbicide options limited, but numerous populations of both weeds have been confirmed resistant to glyphosate.

Over the past two summers (2017/18 and 2019), the Queensland Department of Agriculture and Fisheries (QDAF) Weed Science team have been investigating the impact of growing a competitive sorghum crop on the suppression of these key weeds. Field research has been conducted at the QDAF Hermitage Research Facility located east of Warwick as part of a GRDC-funded project being led by the University of Sydney [US00084 'Innovative crop weed control for northern region cropping systems'].

The field experiments have measured the impact of row spacing and crop density (2017/18) and row spacing and cultivar (2019) in the absence of herbicides (Figure 3). Weed seeds were sown into plots the day after crop planting and weed growth was assessed via biomass measurements and weed seed production counts. In addition, crop yield and biomass have been measured.



Figure 3: Awnless barnyard grass (left) and feathertop Rhodes grass (right) in row spacing trials at Hermitage Research Facility, Queensland. While feathertop Rhodes grass is a more aggressive summer grass weed, a narrow row spacing sorghum can significantly suppress its growth and seed production.

2017/2018 – Higher sorghum density reduced weed seed production

The field trial was planted in early December 2017 with treatments of row spacings at 50, 75 and 100 cm and crop density of 5 and 10 plants m⁻². Weed-free and weedy plots were included. Sorghum emergence was irregular, and the density ranged from 3 to 16 plants m⁻². This range of densities has been taken into consideration in the analysis and interpretation of findings. Weed densities were 15 - 21 BYG and 4 - 6 FTR plants/m². Rain soon after planting caused both weed species to grow rapidly in crop.

Results

In this field trial, neither sorghum row spacing nor density treatments had an impact on BYG or FTR biomass. However, there was a significant overall effect of sorghum density on weed seed production. In sorghum at low densities, both BYG and FTR produced more seed than in sorghum at higher densities (Figures 4 and 5). Sorghum row spacing had no significant effect on weed seed production.

Figure 4: Greater sorghum density significantly reduced seed production of awnless barnyard grass at Hermitage, Queensland.






Figure 5: Greater sorghum density significantly reduced seed production of Feathertop Rhodes grass at Hermitage, Queensland.



Sorghum yield was not significantly different between row spacing or crop density treatments. However, yield was significantly affected by the overall presence of weeds. Weed-free plots averaged a yield of 4.8 t ha⁻¹ (Figure 6). Sorghum yield was reduced 20% by BYG to 3.9 t ha⁻¹ and 44% by FTR to 2.7 t ha⁻¹.

Figure 6: Effect of in-crop awnless barnyard grass and feathertop Rhodes grass on sorghum yield at Hermitage, Qld. No weeds sown indicates plots were maintained weed free. Letters indicate significant ($P < 0.05$) difference at an LSD = 0.34.




2019 – Narrow row spacing reduced weed growth and seed production

Sorghum was sown (mid-January 2019) at either 50 or 100 cm row spacing, comparing three cultivars 85G33, Rippa and MR-Taurus. Crop density was kept consistent at 10 plants/m² (equivalent to 100,000 plants/ha) and weed density was thinned to a consistent 5 plants m⁻². Due to the later planting, the weeds were less vigorous than the previous year.

In brief, the growth of the weeds (biomass and seed production) were not affected by the cultivar of sorghum grown. Supporting this finding is the result that each cultivar had a similar biomass across the row spacings evaluated, so the shading of weeds would have been similar. However, in this trial, row spacing had a significant effect on weed growth with the biomass and seed production of both weeds significantly lower at 50 cm row spacing compared to 100 cm row spacing (Figure 7).

There was no difference in sorghum yield between row spacings and no significant effect of weed presence on yield. Average yield in weed-free plots was 7.55 t ha⁻¹ compared to 7.38 and 7.37 t ha⁻¹ when BYG and FTR were present respectively.

Figure 7: Narrow row spacing (50 cm) in sorghum resulted in a significant reduction in A) seed production and B) biomass, for individual awnless barnyard grass (BYG) and feathertop Rhode grass (FTR) plants. Bars within each graph with a different letter are statistically different at $P < 0.05$.



What do these result mean?

The results achieved across two years show that growing a competitive sorghum crop with increased density and reduced row spacing can significantly suppress the growth and seed production of BYG and FTR.

The impact of row spacing across years was different, and this may be due to the different seasons and growth of the weeds. In 2017/18 the field trial was planted earlier and good rain soon after planting resulted in a large emergence of BYG and robust growth of both weeds. Hence, the weeds had a greater biomass and greater impact on reducing crop yield than in 2019 when the trial was planted later and the weeds did not grow as vigorously.

Cultivar had no impact on suppressing weed growth in the 2019 trial, suggesting cultivar choice will have a lesser impact on sorghum competitiveness than agronomy. However, the impact of cultivar may differ across seasons and locations.

Favourably, growing sorghum at narrow row spacing and increased density, did not have any negative impact on sorghum yield. Therefore, gains in competitiveness and reduction in weed growth can be achieved without reducing yield.

Rarely will a sorghum crop be grown without herbicides. Whether they be residual or knockdown, or a combination of both. Integrating a competitive sorghum crop with herbicides should provide an additive effect on reducing in-crop weed pressures, growth and seed production. Over time, this strategy should deplete the seed banks of both BYG and FTR, and reduce their impact on sorghum production.



Introduction to iGrowth sorghum

What is iGrowth technology?

The iGrowth technology in sorghum was developed by Advanta Seeds through mutagenesis methods and provides tolerance to herbicides of the Imidazolinone family. The iGrowth trait is NOT a GMO. This technology allows farmers to apply registered herbicides at the recommended rates to iGrowth sorghum plants without causing damage. If this herbicide were to be applied on sorghum without this technology, it could cause death or irreversible damage to the crop.



The iGrowth technology will allow Australian sorghum growers the freedom to utilise registered group B herbicides to assist in their integrated weed control programs and will be particularly useful in controlling some common summer grass weeds in their summer crops. Advanta's iGrowth technology also provides the option for double cropping back into paddocks which have grown Clearfield tolerant winter crops, avoiding issues with plant back residuals.

Chemical profile

Imidazolinone herbicides were first developed by American Cyanamid in the 1980's for pre and post-emergence use in various pulse crops including soybeans, peanuts, lucerne, faba beans, chickpea. Imidazolinones inhibit acetolactate synthase (ALS), the enzyme common to the biosynthesis of the branch-chain amino acids (valine, leucine, and isoleucine) and hence formation of proteins.

As pre-emergence treatments, herbicides are taken up by the roots of germinating seedlings, inhibiting growth and in many instances causing plant death. However, it is common for some weed species to emerge initially but fail to develop beyond the seedling stage once newly formed roots access herbicide present in the soil solution. Highly susceptible species die around or very soon after emergence while less susceptible species become susceptible to competition from the developing crop.

Post-emergence applications of IMI herbicides are rapidly absorbed through leaves and translocated via phloem and xylem to sites of meristematic activity where they exert their toxic effect. Post-emergence treatments may also provide pre-emergence activity once the herbicide is moved in by rainfall or overhead irrigation into the soil.

Application window and method

Intervix (imazamox + imazapyr) is primarily a post-emergence foliar applied herbicide with some pre-emergence activity. At the rates registered for use in sorghum (0.75 - 1.0 L/ha), the duration of pre-emergence activity is less than that provided by Lightning (imazamethapyr + imazapyr), Spinnaker (imazamethapyr) and much less than Flame (imazapic) when applied at equivalent amounts of total active ingredient.

Useful Link:

Intervix Label:

https://crop-solutions.basf.com.au/preview/f/product/59cb37fb8d5c5b7640c8374c/Intervix_Label.pdf

Weeds controlled

Imidazolinone herbicides (IMIS) have broadspectrum activity with variation in the activity of individual herbicides for pre-emergence and post-emergence control. Control of broadleaf weeds post-emergence is normally limited to small weeds and relies to a measure on the effectiveness of crop competition occurring subsequent to application particularly for less susceptible species.

Tables 9 and 10 summarise the susceptibility of common weeds of sorghum for the various commercial formulations of IMI herbicides.

Table 9: Susceptibility of grass weeds to IMI herbicides

WEED	INTERVIX 33G/L IMAZAMOX + 15 G/L IMAZAPYR	LIGHTNING 525 G/KG IMAZETHAPYR + 175 G/KG IMAZAPYR	SPINNAKER 700 G/KG IMAZAMETHAPYR COMMENTS		FLAME 240 G/L IMAZAPIC
Grasses	Post	Post	Pre	Post	Pre
Barnyard grass	✓ ✓ ✓	✓	✓	✓	✓ ✓ ✓
Liverseed grass	✓ ✓	✓	✓	x	✓ ✓ ✓
Summer grass	✓ ✓ ✓	✓	✓	x	✓ ✓ ✓
Feathertop Rhodes grass	x	x	✓	x	✓ ✓
Stink grass	✓		✓	x	✓ ✓ ✓
Crowsfoot grass	✓	✓ ✓		x	✓ ✓ ✓
Johnson grass	✓ ✓	✓	✓		✓ ✓
Sweet summer grass			✓ ✓	x	✓ ✓
Dinebra grass					✓ ✓
Native millet					✓ ✓ ✓

x Unsatisfactory control
 ✓ Partial to good control
 ✓ ✓ Very good control
 ✓ ✓ ✓ Excellent control

Table 10: Susceptibility of broadleaf weeds to IMI herbicides

WEED	INTERVIX	LIGHTNING	SPINNAKER 700 GA/KG IMAZAMETHAPYR		FLAME 240 G/L IMAZAPIC
Grasses	Post	Post	Pre	Post	Pre
Amaranth	✓	✓	✓✓	✓	✓✓
Annual ground cherry	✓✓	✓✓	✓✓	✓✓	✓✓
Bellvine/cowvine	✓	✓	✓	✓	✓
Blackberry nightshade	✓	✓	✓		✓
Bladder ketmia	✓	✓	✓✓	✓	✓✓
Burrs	✓	✓	✓	✓	✓
Caltrop/yellow-vine	✓✓	✓✓	✓	✓	✓✓
Common thornapple	✓	✓	✓	✓	✓
Dwarf amaranth	✓	✓	✓✓	✓	✓✓
Fleabane	✗	✗	✗	✗	✗
Melons	✓	✓	✓	✓	✓
Mintweed	✓	✓	✓	✓	✓✓
Parthenium weed			✓✓		
Pigweed (black)	✓	✓	✓✓	✓	✓✓
Pigweed (red)	✓✓	✓✓	✓✓	✓✓	✓✓
Sowthistle (milk thistle)	✗	✗	✗	✗	✗
Wild gooseberry	✓	✓	✓✓	✓✓	✓✓
Wireweed	✓	✓	✓	✓	✓

x Unsatisfactory control
 ✓ Partial to good control
 ✓ ✓ Very good control
 ✓ ✓ ✓ Excellent control



Grass control in iGrowth sorghum

Crop safety

iGrowth sorghum has high levels of tolerance to IMI herbicides having "double gene" tolerance. Some other IMI tolerant crops such as IT corn have lower levels of tolerance when compared to iGrowth sorghum.

Re-crop intervals for IMI herbicides

IMI herbicides are subject to microbial degradation. Conditions favourable for biological activity favour more rapid breakdown of IMI herbicides in the soil. Re-cropping intervals vary with products and rates used. In addition, persistence of IMIs tends to be greater in soils with lower pH (<pH 6) and in low organic matter soils and also where soil moisture is low.

Table 11 summarises re-cropping intervals for commercial IMI formulations. Of the commercial products applied at similar rates of total active constituent, re-crop intervals are shortest for Raptor (imazamox) and Intervix (imazamox + imazapyr) and longest for Flame (imazapic).

Rotational crops most susceptible to carryover of IMIs include canola (except Clearfield varieties), cotton, sorghum (except iGrowth varieties) and corn (except IT varieties). In instances where Spinnaker or Lightning are applied in summer crops at rates above 30 ga/ha, it is prudent if double-cropping to grow Clearfield varieties of either wheat or barley. If re-cropping to non-imidazolinone summer crops, pulses such as mungbean offer the greatest level of safety from the effects of any carryover of Lightning or Spinnaker herbicides.

Persistence of Intervix is expected to be much lower but highly susceptible rotational crops should not be planted within the recommended intervals.

Table 11: Re-crop intervals (months) from in-crop applications of Intervix, Lightning and Spinnaker or fallow applications of Flame in summer rainfall areas.

CROP	INTERVIX	LIGHTNING	SPINNAKER	FLAME 240 G/L IMAZAPIC
Barley or wheat	4	5§§	4 (< 45 g)/15	4§
Clearfield barley or wheat	0	0	0	0
Canola		18	27	36
Clearfield canola	0	0	0	0
Chickpea	4	5	4 (< 45 g)/15	4§
Cotton		18	22	24
Faba beans	0	0	0	3
Maize		10§§§	22	10
IT Maize	0	0	0	3
Mungbean	0	0	0	3
Sorghum		10§§§	22	10
IR Sorghum	0	0	0	0
Soybean	0	0	0	4
Sunflower		18		24
Clearfield sunflower	0	0	0	0

§ plus 200 mm rain
 §§ plus 500 mm of rain or irrigation
 §§§ plus 800 mm of rain or irrigation

Herbicide tank mixes

IMI herbicides are frequently applied alone but are compatible with a range of other selective and non-selective herbicides for the purposes of increasing the range of weeds controlled. Flame for example, is compatible with other pre-emergence herbicides including terbutylazine, diuron and isoxaflutole. IMI herbicides applied in fallow or pre-planting are compatible with knockdown herbicides

Risk of weed resistance

Group B herbicides having the ALS inhibitor mode of action are vulnerable in repeated use in selecting resistant herbicides when used to the exclusion of other weed control methods including herbicides having a different mode of action. The risk of evolution of resistant weed populations is dependent on the frequency of resistant plants in the starting population, the persistence or activity of the applied product, the frequency of application and the history of use of other herbicides sharing the same mode of action prior to the deployment of a new Group B herbicide.

In many sorghum growing areas, Group B herbicides have been sparingly used because of their incompatibility with farming systems including sorghum and cotton where these are quite susceptible to carryover of more persistent Group B herbicides such as metsulfuron, chlorsulfuron, triasulfuron and imazapic. For this reason, there is little evidence of resistance to Group B herbicides in these areas. Even in districts with a longer history of use of imazapic (Flame) in fallow, there have been no confirmed instances of evolved resistant weed populations though this risk remains should this practice be maintained uninterrupted.

Strategies to minimise risk of weed resistance

Weed resistance to IMI herbicides can be avoided by simultaneously employing other weed control measures and in particular, herbicides having differing modes of action. The most prudent approach in incorporating IMI herbicides in conjunction to iGrowth sorghum is to continue use of existing pre and post-emergence herbicides. This approach has been adopted by successful users of Lightning Herbicide in IT maize who found the exclusive use of Lightning often lead to a shift of weed populations that were less susceptible or tolerant to Lightning. In this instance, the inclusion of a pre-emergence herbicide program including s-metolachlor (Dual Gold) and atrazine followed by post-emergence applications of Lightning has enhanced a traditional weed control program relying entirely on s-metolachlor and atrazine and post-emergence treatments capable of controlling broadleaf weeds only.

This approach has also ensured a much lower risk of the development of weed populations resistant to IMI herbicides over the longer term and has already reduced the occurrence of a build-up of naturally tolerant weeds to IMIs.

Good practices for the management of tolerant crops to herbicides

Proper management of herbicide tolerant crops

Group B herbicides having the ALS inhibitor mode of action are vulnerable in repeated use in selecting resistant herbicides when used to the exclusion of other weed control methods including herbicides having a different mode of action.

The risk of evolution of resistant weed populations is dependent on the frequency of resistant plants in the starting population, the persistence or activity of the applied product, the frequency of application and the history of use of other herbicides sharing the same mode of action prior to the deployment of a new Group B herbicide.

In many sorghum growing areas, Group B herbicides have been sparingly used because of their incompatibility with farming systems including sorghum and cotton where these are quite susceptible to carryover of more persistent Group B herbicides such as metsulfuron, chlorsulfuron, triasulfuron and imazapic.

For this reason, there is little evidence of resistance to Group B herbicides in these areas. Even in districts with a longer history of use of imazapic (Flame) in fallow, there have been no confirmed instances of evolved resistant weed populations though this risk remains should this practice be maintained uninterrupted.

Integrated weed management and stewardship

Managing weed resistance: BMP

- Crop Rotation
- Rotate chemical modes of action
- Maintain the use of pre-emergent herbicides
- Use more than one chemical mode of action
- Use non-chemical options: crop competition, mechanical weeding
- Control escapes: keep the seed bank low

Key stewardship rules

- Do not apply to fields with high populations of summer weeds
- Do apply chemical with this mode of action on the target weeds more than once per 12-month period
- Scout for survivors after application, implement control strategies
- Report any serious spray failures
- Control sorghum spp on edges and fence lines
- Clean down equipment to prevent weed spread


How do herbicide resistant weeds develop?

01




Herbicide application

02



Resistant plants survive and generate offspring

03



Repeated use of the same herbicides encourage an increase in herbicide resistant plants

04



In time, the resistant weeds come to dominate



Susceptible biotype



Resistant biotype

[illegible]

[illegible]

For current information on
Pacific Seeds varieties, contact
Toowoomba head office on
(07) 4690 2666 or visit us
online at pacificseeds.com.au

The information provided on this display is intended as a guide only. Advanta Seeds Pty Ltd ('Advanta Seeds') (including its officers, employees, contractors and agents) can not guarantee that every statement is without flaw of any kind. While Advanta Seeds has taken all due care to ensure that the information provided is accurate at the time of publication, various factors, including planting times and environmental conditions may alter the characteristics and performance from plants. Advanta Seeds shall not be liable for any errors or omissions in the information or for any loss, injury, damage or other consequence whatsoever that you or any person might incur as a result of your use of or reliance upon the products (whether Advanta Seeds products or otherwise) and information which appear in this publication. To the maximum extent permitted by law, the liability of Advanta Seeds for any claim whatsoever arising out of the supply or use of or reliance upon the products and information in this publication (including liability for breach of any condition or warranty implied by the Trade Practices Act 1974 or any other law) is limited at its discretion, to the replacement of the products, the supply of equivalent products or the resupply of the publication. For application to specific conditions, seek further advice from a local professional. © Advanta Seeds 2024.